

hep-th/9306116

Reaction-Diffusion Processes as Physical Realizations of Hecke Algebras

Francisco C. Alcaraz ¹ and Vladimir Rittenberg

*Physikalisches Institut, Universität Bonn
Nußallee 12, D - 5300 Bonn 1, Germany*

Abstract

We show that the master equation governing the dynamics of simple diffusion and certain chemical reaction processes in one dimension give time evolution operators (Hamiltonians) which are realizations of Hecke algebras. In the case of simple diffusion one obtains, after similarity transformations, reducible hermitian representations while in the other cases they are non-hermitian and correspond to supersymmetric quotients of Hecke algebras.

¹Permanent adress: Departamento de Física, Universidade Federal de São Carlos, 13560 São Carlos SP, Brasil

It is well known in the literature that several integrable quantum chains corresponding to magnetic systems [1] can be represented as generators of Hecke algebras $H_n(q)$, when certain artificial interactions are added in the bulk and the surface. In this letter we show that the master equation describing the dynamics of some chemical processes limited by diffusion give representations of Hecke algebras, where the supplementary interactions appear naturally.

The Hecke algebra $H_n(q)$ (with $n = L-1$) is an associative algebra with generators e_i ($i = 1, \dots, L-1$) satisfying the relations

$$e_i e_{i\pm 1} e_i - e_i = e_{i\pm 1} e_i e_{i\pm 1} - e_{i\pm 1} \quad (1)$$

$$[e_i, e_j] = 0 \quad ; \quad |i - j| \geq 2 \quad (2)$$

$$e_i^2 = (q + q^{-1}) e_i \quad (3)$$

where q is in general a complex parameter. In our applications q is always real. The main interest in this algebra in statistical mechanics is due to the fact that a spectral-dependent $\tilde{R}(u)$ matrix, satisfying the Yang-Baxter relations [2], can be constructed in a standard form. This construction, also called “Baxterization” [3], implies that the Hamiltonian $\sum_{i=1}^{L-1} e_i$ has an infinite number of conserved charges and this is the first step towards the exact integrability of H .

One can define [4] a sequence (P, M) of quotients algebras of $H_q(n)$ (P and M are positive integers). Since the definition is quite complicated, we will give only a few examples. The quotient $(1, 0)$ implies taking $e_1 = 0$. This quotient is not relevant for our purposes. The $(2, 0)$ quotient is the Temperley-Lieb algebra:

$$e_i e_{i\pm 1} e_i = e_i. \quad (4)$$

The $(1, 1)$ quotient reads:

$$(e_i e_{i+2}) e_{i+1} (q + q^{-1} - e_i) (q + q^{-1} - e_{i+2}) = 0. \quad (5)$$

For our purposes it is sufficient to mention that each (P, M) quotient has as a representative one of the (P, M) Perk-Schultz [5] quantum chains where the Hamiltonian is given by the sum of $H_n(q)$ generators:

$$H^{(P,M)} = \sum_{j=1}^{L-1} e_j^{(P,M)} \quad (6)$$

where

$$\begin{aligned} e_j^{(P,M)} = & \frac{q + q^{-1}}{2} - \left(\sum_{\alpha \neq \beta} E_j^{\alpha\beta} E_{j+1}^{\beta\alpha} + \frac{q + q^{-1}}{2} \sum_{\alpha=0}^{N-1} \epsilon_\alpha E_j^{\alpha\alpha} E_{j+1}^{\alpha\alpha} \right. \\ & \left. + \frac{q - q^{-1}}{2} \sum_{\alpha \neq \beta} \text{sign}(\alpha - \beta) E_j^{\alpha\alpha} E_{j+1}^{\beta\beta} \right). \end{aligned} \quad (7)$$

with $N = P + M$. The $N \times N$ matrices $E^{\alpha\beta}$ have elements:

$$(E^{\alpha\beta})_{\gamma\delta} = \delta_{\alpha\gamma}\delta_{\beta\delta} \quad (\alpha, \beta = 0, 1, \dots, N-1) \quad (8)$$

and

$$\epsilon_o = \epsilon_1 = \dots = \epsilon_{P-1} = -\epsilon_P = \dots = -\epsilon_{P+M-1}. \quad (9)$$

The quantum chains $H^{(P,M)}$ are $U_q(SU(P/M))$ invariant [6] and have the important property that they contain all the irreducible representations of the (P, M) quotient. In the examples we are going to mention, if a chain corresponds to a certain quotient (P, M) then its energy levels (not the degeneracies!) are the same as those of the representative of this quotient given by Eqs.(6-7). We now turn to the physical problem.

It was pointed out recently [7], that the one-dimensional master equation corresponding to a chain with two-body transition rates can be written as a Schrödinger equation with a Hamiltonian described by a quantum chain with nearest-neighbour interactions. Several examples were found in which this Hamiltonian had precisely the structure of Eqs.(6-7). In this letter we present this problem in detail. We will use the notations of Ref.[7].

In order to describe the dynamics of general reaction-diffusion [7] processes we use a master equation. Let us consider an open chain with L sites, where at each site $i = 1, 2, \dots, L$ we attach a variable β taking N integer values ($\beta = 0, 1, \dots, N-1$). By convention we attach the value $\beta = 0$ to a vacancy (inert state) and $\beta = 1, \dots, N-1$ a molecule of type β . The master equation describing the time evolution of the probability distribution $P(\{\beta\}; t)$ is

$$\begin{aligned} \frac{\partial P(\{\beta\}; t)}{\partial t} = & \sum_{k=1}^{L-1} \left[-w_{0,0}(\beta_k, \beta_{k+1}) P(\beta_1, \dots, \beta_L; t) \right. \\ & \left. + \sum'_{\ell, m=0}^{N-1} w_{\ell, m}(\beta_k, \beta_{k+1}) P(\beta_1, \dots, [\beta_k + \ell]_N, [\beta_{k+1} + m]_N, \dots, \beta_L; t) \right] \quad (10) \end{aligned}$$

where $w_{\ell, m}$ are the transition rates and the prime in the second sum indicates the exclusion of the pair $\ell = m = 0$. In (10) the symbol $[x + y]_N$ means the addition $(x + y)$, modulo N . We restrict ourselves to the cases where the transition rates are homogeneous and depend only on the molecules on nearest neighbouring sites. The transition rate $w_{\ell, m}(\alpha, \beta)$ gives the probability, per unit of time, that an initial state where two consecutive sites are occupied by molecules of type $(\alpha + \ell, \beta + m)$ end up occupied by molecules of type (α, β) . The rate $w_{0,0}(\alpha, \beta)$ gives the probability, per unit of time, that the configuration of two neighbouring sites (α, β) are changed. From these definitions the conservation of probabilities implies

$$w_{0,0}(\alpha, \beta) = \sum'_{r,s} w_{r,s}(\alpha - r, \beta - s), \quad (11)$$

where $r = s = 0$ is again excluded.

The master equation (10) can be interpreted as a Schrödinger equation

$$\frac{\partial}{\partial t} |\Psi\rangle = -H |\Psi\rangle \quad (12)$$

in Euclidean time, if we identify the probability $P(\{\beta\}, t)$ as the wave function. The Hamiltonian in (12)

$$H = \sum_{j=1}^{L-1} H_j \quad (13)$$

acts in a Hilbert space of dimension N^L , while H_j acts in the subspace $V^{(j)} \otimes V^{(j+1)}$ and is given by

$$H_j = U_j - T_j, \quad (14)$$

where


$$T_j = \sum_{\ell, m=0}^{N-1} \sum_{\alpha, \beta=0}^{N-1} w_{\ell, m}(\alpha, \beta) E^{\alpha, [\alpha+\ell]_N} \otimes E^{\beta, [\beta+m]_N} \quad (15)$$

$$U_j = \sum_{\alpha, \beta=0}^{N-1} w_{0,0}(\alpha, \beta) E^{\alpha\alpha} \otimes E^{\beta\beta}. \quad (16)$$


The above Hamiltonian describes the time evolution from an initial probability $\Psi_0(\{\beta\}, t=0)$ with dynamical processes given by the transition rates $w_{\ell, m}(\alpha, \beta)$.

Here are some examples of processes in which the H_j 's of Eq.(13) coincide with the generators e_j of the Hecke algebra $H_n(q)$.

a) **diffusion processes:** we consider $(N-1)$ types of molecules $(A_1, \dots, A_{N-1})$ which diffuse to the right:




with equal rates Γ_R and to the left:



with equal rates Γ_L . Choosing the time scale in such a way that the diffusion constant $D = \sqrt{\Gamma_L \Gamma_R} = 1$ and introducing $q = \sqrt{\Gamma_R / \Gamma_L}$, up to an equivalence transformation, we obtain $H_j = e_j$ corresponding to the quotient $(2, 0)$.

b) **diffusion and interchange processes:** to the diffusion processes (17) and (18) we add the interchange to the right processes:




with rates Γ_R and with rates Γ_l if $b < c$. In this case the quotient is $(N, 0)$ and the obtained representation of $H_n(q)$ is hermitian (the same applies to the diffusion processes of case a).

c) **diffusion, interchange and coagulation processes:** we now add to the processes (17-19) only one reaction for the molecule A_1 (the molecules are ordered!)






with a rate Γ_R and the processes where $\emptyset$ and A_1 interchange their position in the final state with rate Γ_L . In this case the quotient is $(N-1, 1)$ and the representation of H_n is non-hermitian. As a rule, non-hermitian representations occur for quotients (P, M) with $M \neq 0$ i.e. when the representative Hamiltonians $H^{(P,M)}$ are supersymmetric.

One obtains new quotients of H_n generalizing the reaction (20) by taking




where $s = r \pm 1$, $s' = r' \pm 1$, and $r \neq r', s'; s \neq r', s'$. In this way one obtains the quotient $(N-2, 2)$. Taking more reactions like in Eq.(21,22) one can get arbitrary quotients (P, M) .

d) **diffusion-polymerisation (modulo N) processes:** in this case besides the diffusion (Eqs. 17-18), one considers the processes:



For all these processes one obtains the quotients $(1, 1)$. Let us remark that the processes described by Eq. (23-25) were already considered in Ref.[9] and called "P-poly" for irreversible diffusion cluster-cluster aggregation processes. Using probabilistic methods several exact results were obtained (see also Ref.[10]) concerning the concentrations of the different types of molecules. We believe that the profound reason which explains why these results could have been obtained is that the $(1, 1)$ quotient has a spectrum which can be obtained using free fermion methods [11]) (see also [12]).

e) **diffusion, interchange and annihilation processes:** because of interchange processes, the labels of the molecules are ordered; we can now add the following two processes:



with $\Lambda_1 + \Lambda_{-1} = \Gamma_R + \Gamma_L$. In this way we obtain again the quotient $(2, 1)$.

The reader might wonder how the above $H_n(q)$ quotients were obtained. From the work done in Ref.[7] we have developped a certain intuition for chemical processes up to $N = 3$. We have then used the computer to check that no other solutions exist for $N = 2$ and 3 and generalized the solutions obtained for N up to 3 to all

N and checked them. There are certainly other solutions which one can discover studying systems with $N = 4$ and certainly examples of the quotients $(2, 2)$ different from the one given in Eq.(21,22) will show up. We hope that the present effort to connect representatives of $H_n(q)$ and chemical processes will help to compute various correlation functions for the latter. One immediate consequence of the above mentioned connexion is that all processes with $q \neq 1$ are massive (the concentrations for example have an exponential fall-off in time) and all those with $q = 1$ are massless (algebraic fall-off).

Acknowledgements

F.A. thanks the Conselho Nacional de Desenvolvimento Científico and Tecnológico - Brasil and the Deutsche Forschungsgemeinschaft - DFG - Germany for support.

References

- [1] P.P. Martin, Potts Models and Related Problems in Statistical Mechanics, World Scientific, Singapore 1991
- [2] R. J. Baxter, Exactly Solved Models in Statistical Mechanics, Academic, N. Y. 1982
- [3] V.R. Jones, Int. J. Mod. Phys. **B4** (1990), 701.
- [4] P.P. Martin and V. Rittenberg, Int. J. Mod. Phys. **A7**, Suppl. 1B 1992), 707 and references therein
- [5] J.H.H. Perk and C.L. Schultz in "Non-Linear Integrable Systems, Classical theory and Quantum Theory", Ed. by M. Jimbo and T. Miwa, World Science, Singapore 1981
- [6] T. Deguchi, J. Phys. Soc. Jpn. **58** (1989), 3441; T. Deguchi and Y. Akutsu, J. Phys. **A23** (1990), 1861; M.L. Ge, G.C. Liu and K. Hue, Mod. Phys. Lett. **A6** (1991), 1173
- [7] F.C. Alcaraz, M. Droz, M. Henkel and V. Rittenberg, "Reaction-Diffusion Processes, Critical Dynamics and Quantum Chains", University of Genève preprint UGVA-DPT 1992/12-799, Annals of Physics (to be published)
- [8] M.V. Smoluchowski, Z. Phys. Chem. **92** (1917), 129; A.A. Ovchinnikov and Y.B. Zeldovich, Chem. Phys. **28** (1978), 215; D. Toussaint and F. Wilczek, J. Chem. Phys. **78** (1983), 2642; F. Leyvraz and S. Redner, Phys. Rev. **A46**(1992), 3132

- [9] J.L. Spouge, Phys. Rev. Lett. **60** (1988), 871
- [10] V. Privman, Phys. Rev. **A46** (1992), 6140
- [11] T.D. Schultz, D.C. Mattis and E.H. Lieb, Rev. Mod. Phys. **36** (1964) 856
- [12] R.J. Glauber, J. Math. Phys. **4** (1963), 294; B.U. Felderhof, Rep. Math. Phys. **1** (1970), 219; Phys. Lett. **120A** (1987), 135; A.A. Lushnikov, Sov. Phys. JETP **64** (1986), 811; D. ben-Avraham, M.A. Burschka and C.R. Doering, J. Stat. Phys. **60** (1990), 694